



IMS user group
integrated monitoring system


event 2019

IMS overview

Summary of data recorded for 2018-19

**Service providers
190 locations
2018-19**

**89 agency codes
101 pharmacies**



2018-19



69,115

Syringe exchange visits




65,610

Interventions delivered




20,050

Clients recorded



6 year trend – all IMS clients




6 year trend – client cohort groups

What's changed

- Changes to services
 - Provider & pharmacy changes
- Recording systems and processes
 - PharmOutcomes





3 year trend – local authority areas

Activities recorded

Summary of activity recorded in IMS

Core IMS modules

Interventions

- Advice & Info - 6,605
- Brief Intervention – 59,005



329 intervention type descriptions used

Referrals

- 'IN' Source - 632
- 'OUT' onward – 510



52 referral types or organisations

Syringe Transactions

- Exchange visits - 69,115
- Returns - 21,885



Return recorded for 31.2% of visits

Wellbeing reviews

- 'WEMWBS' – 2,113

Reviews recorded for 1,688 people

Includes 291 people with 2 or more



Cohorts & demographics

What IMS data shows about our service users

6,974



Brief Intervention
Clients - Drug or Alcohol

- 61% male, 39% female
- 63% aged between 30 and 59
- Average 4 interventions

7,117

- 81% male, 19% female
- 70% aged between 30 and 49
- Average 8 needle exchange visits

People Who Inject
Psychoactive Drugs



People Who Inject
Drugs: Steroids

5,959

- 96% male, 4% female
- 77% aged between 19 and 39
- Average 2 needle exchange visits




- 1,529 (22%) had no substance or did not record one
- 119 (2%) someone else's use

6,974

Main substances

- Alcohol 3,187 (46%)
- Heroin 844 (12%)
- Cannabis 419 (6%)
- Cocaine 402 (6%)



7,117

Main substance recorded for 5,388 people (76%)

- Heroin 4,220 (78%)
- Crack cocaine 217 (4%)
- Amphetamines 82 (2%)


Second substance recorded for 1,779 people (33%)

- Crack cocaine 1,344 (76%)
- Heroin 127 (7%)
- Steroids 69 (4%)

5,959

Second substance recorded for 235 people (4%)

- Another Steroid or IPED 195 (83%)
- Other Drugs (5%)
- Cocaine (3%)
- Heroin (2%)
- Alcohol (2%)



Maps

<https://ims.ljmu.ac.uk/maps>

